

List of Illustrations

Nathaniel Charles Rothschild 1877–1923	Frontispiece
Charles Rothschild (1892). He published his first paper on Lepidoptera at Harrow School. (Photographer not known.)	2
Ashton Wold, Northamptonshire, the home of Charles Rothschild. (Photographed by Miriam Rothschild).	3
Csételek, Hungary, Rothschild's ranch on his father-in-law's estate. (Photographed by Clarisse Strasser)	3
Hugh Birrell, a fellow undergraduate at Cambridge, who received 60 letters from Charles Rothschild between 1899–1908. (Photographer not known)	4
Charles Rothschild in 1907 outside his father-in-law's house at Csételek, Hungary. (Photographed by Clarisse Strasser)	7
Roszika von Wertheimstein in the year of her marriage to Charles Rothschild. (Photographer not known)	9
Francis Henley and a friend at the Pyramid of Cheops. Charles Rothschild discovered the plague-carrying flea in Egypt, and named it <i>Xenopsylla cheopis</i> . (Photographed by Charles Rothschild)	10
Charles Rothschild in 1922 before his last collecting expedition to Bettyhill, Sutherland.	11
Charles Edward Fagan, 1855–1921. Natural History Museum archive.	14
William Ogilvie-Grant, 1863–1924. Natural History Museum archive.	14
Francis Henley, a life-long friend of Charles Rothschild.	14
Box Hill, Surrey, c. 1910, looking west towards Ranmore Common.	17
Cheddar Gorge c. 1910, then bare and free of scrub, with the road as yet unmetalled.	17
Viscount Ullswater (previously J.W. Lowther), Speaker of the House of Commons, President of the SPNR from 1915 to 1931. (Photographer not known)	21
Lionel Walter, Lord Rothschild, President of the SPNR from 1931 to 1937. (Photographer not known)	22
Sir John Fryer, 1886–1948, agriculturist and conservation pioneer.	22
Neville Chamberlain, Walter Rothschild, R. Bowdler Sharpe, F. Gayner and Mr. Kenrick, photographed by Charles Rothschild before a day's collecting Lepidoptera at Ashton Wold.	23
A completed questionnaire for Hengistbury Head, Hants, proposed by C.J. Wainwright.	26

The corresponding map, with the proposed reserve marked with rough pencil lines. The sand bar was longer then, and apparently not yet cluttered with buildings and coastal defences.	26
An unusually detailed questionnaire for Freshney Bog, Lincs, completed by Miss Florence Woolward.	28
Distribution of the areas 'worthy of protection' selected by the SPNR in 1915. Category A were those considered to be the most outstanding. Category B form the bulk of the selection, while Category C were of 'secondary importance' (from Sheail, 1976).	31
Wistman's Wood on Dartmoor, a place of special interest to the first generation of plant ecologists.	35
The 'Rothschild Reserves' archive, stored in two wooden museum drawers, now at the Wildlife Trusts headquarters, Lincoln.	36
A typical 'site dossier' from the SPNR archive – maps, standard form and correspondence – all filed in the Rothschild bank's 'blue linen' envelopes.	36
Sutton Heath, Northants, is marked by a simple cross. Presumably the area of rough ground shown on the contemporary 1" map is the proposed reserve.	37
The proposed reserve at Newham Lough, Northumberland, is coloured in on the Society's large-scale map.	37
'Kingley Bottom' site roughly that of today's National Nature Reserve, purchased by the Nature Conservancy in 1952.	37
Box Hill is marked only by a rough circle, which excludes most of the special box groves the SPNR wanted to protect!	37
Richardson Carr, at Home Farm, Tring, where he kept a large aviary of tropical finches. He was a friend and agent of Charles Rothschild, and together they wrote over a thousand letters to owners of potential nature reserves. (Photographer not known)	39
A page from the 'Rothschild List' of places 'worthy of protection' printed in 1915. Saltmarsh and sand bar at Blakeney Point, purchased for the National Trust by Charles Rothschild and others in 1912. (Photographed by Peter Wakely)	44
Hand-coloured map of Ray Island and Old Hall Marshes from the SPNR archive. Woodwalton Fen in 1948. (Photographed by S. Beaufoy)	46
The bungalow built by Charles Rothschild at Woodwalton Fen (1936). (Wildlife Trusts archive)	47
The Society's maps of the old quarry at Swaddiwell Field, Northants, at 1" and 6" to the mile scale.	48
Views of Ashdown Forest taken 50 years apart show extensive spread of bracken and scrub with the loss of botanically rich habitats. (Photographed by C.J. Marrable)	54
The 'Lewes Downs' site shows the scrub invasion that has overtaken so many grassland sites since 1916. Part of the area is now the Malling Down nature reserve, owned by the Sussex Wildlife Trust. The chalk pits were made in the nineteenth century. (Photographed by Peter Marren)	55
Scotton Common, Lincs., then wild heath and bog, is now mostly conifer plantation.	58
Oakley Purlieus, Northants., overlay deposits of ironstone. It was clear-felled during the 1940s, quarried and afterwards reclaimed for agriculture.	58
Harlestone Heath, once one of Northamptonshire's richest botanical sites, is now mostly plantation or reclaimed for housing.	59

Much of the SPNR site at Aston Upthorpe Downs, Berks., has since been ploughed or converted to botanically dull grassland, including 'The best area' around Lowbury Hill.	59
Wicken Fen c. 1910, more open then than now, partly through mowing for thatch and marsh hay.	65
Larva of swallowtail butterfly at Wicken Fen, probably photographed by Charles Rothschild.	65
Today the foreground is much the same, but the fen behind has been heavily scrub-invaded.	65
The Pasqueflower has disappeared from SPNR sites at Aldbury Nowers, Fleam Dyke, Swaddiwell Field and Aston Upthorpe. (Photographed by Peter Wakely)	69
Spiked speedwell disappeared from Garboldisham Heath around 1980, and is now among the rarest Breckland specialties. (Photographed by Peter Wakely)	69
The Silver-spotted Skipper, lost since 1915 from Aldbury Nowers, Devil's Dyke and Kingley Vale. (Photographed by Bob Gibbons)	70
Rothschild hoped to protect the Essex Emerald moth by preserving its habitat on the Essex saltings. The moth is now nationally extinct. (Photographed by Paul Waring)	73
The Black-veined was another rare moth that Rothschild tried to preserve. Its decline to endangered status was not through collecting, as he feared, but from habitat destruction and scrub invasion. (Photographed by Paul Waring)	73
The best site for the Sandhill Rustic moth, at Lytham St. Anne's, Lancs., was destroyed through building as early as 1919. Rothschild substituted another site for it near Fleetwood, but coastal protection works have since destroyed that one too. (Photographed by Paul Waring)	73
The Alpine Catchfly still occurs at the Lake District site which formed one of the SPNR's proposed 'species reserves' (Photographed by Hugh Lang)	74
White Rock-rose has gone from the section of cliff proposed for it at Babbacombe, but survives at Berry Head at the opposite end of Torbay. (Photographed by Peter Marren)	74
The naturalised herb, <i>Asarabacca</i> , with its shiny, cyclamen-like leaves, was among the reasons for proposing Weston Turville reservoir, Bucks., as a nature reserve. The original site is now built on, but the species was transplanted nearby. (Photographed by Peter Marren)	75
'The world's smallest nature reserve' (300 square yards) was established at Badgeworth, Glos., in 1932 to protect the Adder's-tongue Spearwort. (Photographed by Peter Marren)	77
The distribution of National Nature Reserves in England in 1993. (English Nature)	81
Two views: The great yews of Kingley Vale in 1910 and 1993 – hardly changed in nearly a century, though bearing the bullets and scars of military manoeuvres of two world wars. (Photographed by Peter Wakely)	82
The sparrow of tomorrow? A bird with a dramatic reduction in numbers. Tring Reservoir 1906. (Photographed by Oliver Pike)	85
Lady's Slipper, <i>Cypripedium calceolus</i> . (Photographed by Charles Rothschild)	87
Lesser Butterfly Orchid, <i>Plantathera bifolia</i> . Photographed by Charles Rothschild around 1900. A greatly decreased wild orchid.	87
Sunset over Weston Turville reservoir, 1936. (Photographed by Drogo Montagu)	90
Abbots Moss in 1912: 'the breeding ground of the Marsh Ringlet' (Large Heath butterfly). You could probably find a similar scene today (less the collectors!), though the moss has dried out somewhat and suffers from pines. (Wildlife Trusts archive)	93

Burwell Fen, part of the SPNR's Adventurers' Fen, at the turn of the century, still wet and probably cut for marsh hay. (Photographer not known)	93
The dunes at Ainsdale, Lancs., finally became a nature reserve in 1965. (Photographed by Peter Wakely)	95
Barton Hills survived the war, just, got over myxomatosis, and today is one of the best preserved chalk grassland sites on the list. (Photographed by Peter Wakely)	104
Berry Head is a combination of public amenity and nature reserve, and has escaped the development which engulfed the cliffs nearer to Torquay. (Photographed by Peter Marren)	107
Brean Down c. 1910, when ravens and peregrines nested on these cliffs. The scene is essentially the same today, but the upper slopes have grown scrubbier.	113
Chichester Harbour today. (Photographed by Peter Wakely)	121
The top of Coombe Hill at the turn of the century, then heath and gorse scrub. It is now grass and gorse scrub.	123
Crag Lough as it appears on the SPNR's map. Note the placing of the pencil cross; apparently Rothschild was more interested in the wet field below the crags than the lough itself.	126
Dawlish Warren, a beckoning finger of sand, enclosing a salt marsh from the O.S. map of 1912 with the site marked by crosses.	129
The Devil's Kneading-Trough at Wye Downs, Kent, listed by Rothschild for its rare insects. (Photographed by Peter Wakely)	131
Cormorant Island, part of the Farnes, in 1905, with gulls in the foreground. (Photographer not known)	136
The Pinnacles, Farne Islands, c. 1905. The Farnes is one of England's most successful nature reserves, though not formally named as such until 1993.	136
Forge Valley Woods, which follow the steep-sided valley of the Derwent, was created a National Nature Reserve in 1977. (Photographed by Peter Wakely)	139
Goonhilly Downs is big enough to absorb earth tracking stations, airfields and plantations and yet retain much of its former value for wildlife. (Photographed by Peter Wakely)	142
Hambledon Hill is still an impressive site. It was 'declared' a National Nature Reserve in 1993. (Photographed by Peter Wakely)	146
Hartlebury Common has shrunk by about a third since the SPNR listed it. It is now a Local Nature Reserve. From the SPNR's Contemporary Ordnance Survey map.	149
The SPNR's map showing Hutton Roof seems to include the village with the limestone scenery beyond.	159
Inkpen Crocus Field at crocus time in late March. The field was purchased by the Bucks, Berks and Oxon Naturalists Trust in 1985 to save it from possible housing development. (Photographed by Peter Marren)	160
Kynance Cove has suffered from crowds and erosion, but on quiet days retains its charm and beauty. (Photographed by Peter Marren)	164
Littlesea, today part of Studland National Nature Reserve. (Photographed by Peter Wakely)	167
Naphill Common, Bucks., open and grazed in 1900, is now woodland, barely distinguishable from neighbouring woods.	172
Pusey Camp, once a noted botanical 'locality' is no longer of special interest. (Photographed by Peter Marren)	178
Rostherne Mere, Cheshire, still looks pretty, and is a National Nature Reserve, but beneath the surface has become polluted from nitrate fertiliser and sewage. (Photographed by Peter Wakely)	181

Roudsea Wood, ancient woodland along the shore of Morecombe Bay. (Photographed by Peter Wakely)	182
The rich flora of St. Vincent's Rocks, beneath Clifton Suspension Bridge, has suffered from roadworks, scrub invasion and metal pollution. (Photographed by Peter Marren)	186
Beds of sea lavender at Gibraltar Point, part of Rothschild's 'Skegness' site. (Photographed by Peter Wakely)	192
Slapton Ley at Start Point seen from the north, now a nature reserve much used for field studies. (Photographed by Peter Wakely)	195
A kingfisher at her nesting hole, Tring and Weston Turville Reservoirs. (A classic photograph by Oliver Pike, 1906)	203
Great Crested Grebe sitting on her nest at Tring and Weston Turville Reservoirs. (Photographed by W. Bickerton, 1906)	204
Black-necked Grebe, photographed by Oliver Pike in 1919 on the occasion of its nesting at Tring reservoirs, the first nesting record in Britain.	204
An old brickpit and reedbeds at Wicken Fen. (Photographed by Peter Wakely)	206
Sugar limestone (and distant reservoir) at Widdybank Fell, 'a most wondrous botanical station'. (Photographed by Peter Marren)	208
Winterton Dunes, Suffolk, became one of the first National Nature Reserves in 1956, though by then much damaged and reduced since 1915. (Photographed by Peter Wakely)	209
Woodchester Park, with its famous unfinished mansion, was turned into an 'alpine valley' in the 1950s by massed conifer planting.	211
Four views of Wood Walton Fen in the 1930s. (Wildlife Trusts archive)	212
Dowles Brook, running through the heart of the Wyre Forest.	215
Half of Thorne Moors, Yorks., has been stripped and laid bare by industrial peat milling. The best remaining area lies on early twentieth century peat workings. (Photographed by Peter Roworth)	227
Near natural woodland at Bramshaw Wood, New Forest. (Photographed by Peter Marren)	232
A peaceful scene in the New Forest, probably not much changed in 1990 if you discount the busy roads nearby.	232