

Mushroomology

Mushroomology

Edited by

Jianping Xu
Timothy James
Dapeng Bao



BRILL | WAGENINGEN ACADEMIC

Cover image: shiitake mushrooms in a factory in Hamilton, Canada. Photo made by Jianping Xu.

Library of Congress Cataloging-in-Publication Data

Names: Xu, Jianping (Professor of Biology), 1965- editor | James, Timothy Yong, 1973- editor | Bao, Dapeng editor
Title: Mushroomology / edited by Jianping Xu, Timothy James, Dapeng Bao.
Description: Leiden ; Boston : Brill : Wageningen Academic, [2026] | Includes bibliographical references and index.
Identifiers: LCCN 2025051018 (print) | LCCN 2025051019 (ebook) | ISBN 9789004751675 hardback | ISBN 9789004751699 ebook
Subjects: LCSH: Mushroom culture | Mushrooms | Cultivated mushroom | Cultivated mushroom--Breeding | Cultivated mushroom--Genetic engineering
Classification: LCC SB353 .M87 2026 (print) | LCC SB353 (ebook)
LC record available at <https://lcn.loc.gov/2025051018>
LC ebook record available at <https://lcn.loc.gov/2025051019>

Typeface for the Latin, Greek, and Cyrillic scripts: "Brill". See and download: brill.com/brill-typeface.

ISBN 978-90-04-75167-5 (hardback)

ISBN 978-90-04-75169-9 (e-book)

DOI 10.1163/9789004751699

Copyright 2026 by Koninklijke Brill BV, Plantijnstraat 2, 2321 JC, Leiden, The Netherlands, except where stated otherwise.

Koninklijke Brill BV incorporates the imprints Brill, Brill Nijhoff, Brill Schöningh, Brill Fink, Brill mentis, Brill Wageningen Academic, Vandenhoeck & Ruprecht, Böhlau and V&R unipress.

All rights reserved. No part of this publication may be reproduced, translated, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without prior written permission from the publisher. Requests for re-use and/or translations must be addressed to Koninklijke Brill BV via brill.com or copyright.com.

For more information: info@brill.com.

This book is printed on acid-free paper and produced in a sustainable manner.

Contents

- Figures and Tables ix
- 1 Diversity and taxonomy of edible and medicinal mushrooms 1
Gang Wu and Zhu-Liang Yang
 - 2 Diversity of life cycles, breeding systems, and mating types in mushrooms 58
Ursula Kües and Ines Teichert
 - 3 Mushroom population genetics and germplasm 154
Kuan Zhao and Jianping Xu
 - 4 Mushroom physiology and nutritional requirements 189
Juanjuan Liu, Xinlei Zhang, Na Pang, Rong Zhu and Zemin Fang
 - 5 Quantitative trait loci mapping and mushroom breeding 215
Anton S.M. Sonnenberg and Arend F. van Peer
 - 6 Mushroom genomics – the history, driving forces and current state of the genomics of mushroom-forming fungi 239
László G. Nagy, Balázs Bálint and Igor V. Grigoriev
 - 7 Molecular biology and biotechnology of mushrooms and genetically modified mushrooms 278
Rui Liu, Jing Zhu, Ang Ren, Liang Shi and Mingwen Zhao
 - 8 Gene editing in edible and medicinal mushrooms and applications 306
Pengtao Zhang, Mengqian Liu, Mengfei Zhang, Guoliang Meng and Caihong Dong
 - 9 Mutation and strain degeneration in mushrooms 326
Dr. Timothy Y. James
 - 10 *Agaricus* mushroom pests and pathogens 351
Dr. David Meigs Beyer and Dr. Mike Wolfin
 - 11 *Agaricus bisporus*, the button mushroom 394
Jianping Xu

- 12 Biology and cultivation of oyster mushrooms 411
Wei Gao and Chenyang Huang
- 13 *Lentinula edodes*, the Shiitake Mushroom 432
Dapeng Bao
- 14 *Volvariella volvacea*, the straw mushroom 450
Junsheng Fu
- 15 *Flammulina filiformis*, the golden needle mushroom 466
Yongxin Tao
- 16 *Tuber* spp., the true truffles 481
Bang Feng
- 17 Biology and cultivation of *Auricularia heimuer* and *Auricularia cornea* 501
Fangjie Yao, Xiaoxu Ma and Yuling Cui
- 18 *Phlebopus portentosus*, the cultivated bolete 520
Yang Cao and Kaiping Ji
- 19 *Hericiium erinaceus*, the lion's mane mushroom 540
Hongyi Han and Jianping Xu
- 20 *Ganoderma sichuanense*, the cultivated lingzhi 556
Ke Leqin and Wu Qingqi
- 21 *Cyclocybe chaxingu*, the chashugu 578
Yang Gao, Hai-Yang Song, Hai-Jing Hu and Dian-Ming Hu
- 22 *Cordycipitoid fungi* (*Cordyceps sensu lato*) 595
Yongjie Zhang, Chunru Li, Zhengming Qian and Chenghua Zhang
- 23 Biology and cultivation of true morels (*Morchella* spp.): current progress and future trends 608
Xi-Hui Du, Jie Tang, Hao Tan, Jia Ling, Yong Wang and Wei-Hong Peng
- 24 Mushroom culture 635
Jianping Xu, Han Hongyi, Ting Chen, Wu Zijing and Han Shenghua

- 25 Mushroom pigments: biosynthesis, diversity, ecological roles, and applications 658
Lan Lin and Jianping Xu
- 26 Mushroom processing and processed products 706
Liping Liu, Jun He and Yanfang Liu
- 27 Species diversity and poisoning classification of poisonous mushrooms 733
Wangqiu Deng and Taihui Li
- 28 Mushroom volatiles – diversity, synthesis, function, and applications 753
Annika Wagner, Nikolas Hoberg and Martin Rühl
- 29 Introduction to home growing mushrooms 790
Andrew Langevin
- 30 Mushrooms in a changing world 805
Yuan Guo, Shuang Song and Shouxian Wang
- Index 837

Figures and Tables

Figures

- 1.1 Diversity of edible and medicinal mushrooms across continents. In sub-panel *a*, the total number of edible and mushroom species, the number of edible species (E) and the number of medicinal species (M) for each continent (Africa, Asia, Europe, North America, South America, and Oceania) are provided separately. In sub-panel *b*, the vertical orange bars represent the numbers of unique species for each continent and their shared species with other continents, categorized below by gray dots and vertical lines. The horizontal blue bars display the total number of species for each continent. 8
- 1.2 The number of edible and medicinal mushrooms within families of Basidiomycota and Ascomycota as well as all described species of each family. The relationships among families were shown in an artificial family-level phylogenetic tree. Some genera with edible and medicinal mushrooms are retained their generic names due to their uncertain attributions at the family level. In the panel, the column of *all* represents the total number of described species of the corresponding clades, and that of *edible & medicinal* represents the total number of edible and medicinal mushrooms of the corresponding clades, while those of *edible* and *medicinal* represent the number of edible and medicinal mushrooms, respectively. The orders to which the included families as well as some genera belong are labelled in the purplish and greenish boxes. 10
- 2.1 Generalized schemes of the three major breeding systems within the fungal subkingdom Dikarya. Schematized are filamentous species that develop fruiting bodies for sexual spore production. Haploid nuclei are indicated by small circles within cells, with distinct genotypes (hereby explicitly referring to the mating type information present in heterokaryotic mycelia) distinguished by filled and open circles, respectively. For the sake of simplicity, the scheme on the left only considers life cycle control in unifactorial heterothallics by a single *mat* locus with bipolar meiotic segregation. In such cases, the typical meiotic tetrad consists of two nuclei of each parental mating type which are individually enclosed in the four different meiospores. Bifactorial heterothallics (not illustrated), on the other hand, may contain in their typical tetrads either two nuclei of each parental mating type, or two nuclei of two new recombinant parental mating types, or one nucleus per each parental mating type and each new recombinant mating type. In pseudo-homothallics with a single *mat* locus control, the sexual spores are heterokaryotic and contain nuclei of both parental mating types, as shown in the diagram on the right. In pseudo-homothallics with two *mat* loci (not illustrated), the spores can either carry both parental mating types or both recombinant mating types. 61

- 2.2 Life cycle of the model ascomycete *Neurospora crassa*. The heterothallic species has haploid mycelia of two alternate mating types *mat A* (= *mat1-1*) and *mat a* (= *mat1-2*) which can undergo the same developmental processes, as indicated by two *mat*-specific life cycles (the left one for *matA* and the right one for *mat a*). The haploid, dikaryotic and diploid stages in the life cycles are indicated by light grey, white and dark grey shading. Main environmental factors that positively (black letters with white background) or negatively (white letters with grey background) influence steps in the life cycles are indicated. Each mating type can form haploid female (nucleus-receiving) sexual structures, the ascogonia, that are enveloped by sterile hyphae, thus forming the protoperithecia. These are fertilized by mitotic spores or vegetative hyphae of the opposite mating type with the help of a specialized female cell, the trichogyne. Fertilization leads to dikaryon formation and the generation of ascogenous hyphae that are a prerequisite for ascus and ascospore formation inside the growing haploid fruiting body (perithecium). Note that the dikaryotic stage in ascomycetes is very short. In the young dikaryotic ascus, karyogamy takes place between haploid nuclei of different mating types, followed by the two meiotic division and one post-meiotic mitosis, after which eight haploid ascospores develop in the ascus. Note that information here is limited to environmental factors, mating-type loci and pheromones/pheromone receptors, because description of the many known genetic determinants is beyond the purpose of this review. 74
- 2.3 Reciprocal courtship and fertilization of *Neurospora crassa* between two strains of different mating type is controlled by mating-type-specific gene expression of pheromone and pheromone receptor genes. Each figure part shows a scheme of the interaction between two different mating types, with one acting as female partner, generating ascogonial coil and protoperithecium with trichogyne, and the other one acting as male partner, generating a microconidium. Macroconidia and vegetative cells can also act as male partner, but are not included here for clarity. During fertilization, a “male” nucleus from the microconidium is translocated through the trichogyne to the “female” ascogonium. Below the schemes, a detailed cellular view of the interaction between trichogyne and conidium is given, including details of mating type-specific gene expression. **A.** Courtship between a female-acting *mat1-1* strain and a male-acting *mat1-2* strain. As shown at the top, a female-acting *mat1-1* strain forms protoperithecia with ascogonia (ascogonial coils) at their base (shown in white), from which a trichogyne (also shown in white) grows and searches for a suitable attracting male cell for courtship (e.g. a microconidium shown in grey acting as spermatium) of a male-acting *mat1-2* strain and which finally takes up the cell nucleus from it during fertilization. **B.** Courtship between a female-acting *mat1-2* strain and a male-acting *mat1-1* strain. The situation is reciprocal to the one shown in **A.** with the *mat1-2* strain being the female-acting partner (with ascogonial coil and trichogyne shown in grey) and the *mat1-1* strain being the male-acting partner (with a microconidium acting as spermatium shown in white). In the detailed scheme of the enlarged cells in **A.** below, *mat1-1* is located in the nucleus of the female trichogyne and *mat1-2* in the nucleus of

the spermatium. In **B**, *matl-2* is located in nucleus of the female trichogyne and *matl-1* in the nucleus of the spermatium. The nuclei with crucial chromosomal segments are highlighted in grey. The structures of the alternate *matl* idiomorphs on chromosome 1 (Chr 1) are indicated in the respective *matl-1* and *matl-2* nuclei with genes being either shown in white (*matl-1*) or in grey (*matl-2*) while conserved neighbouring genes are shown in black (including an ATPase gene *ar-2* needed in *N. crassa* for sexual development, Randall and Metzzenberg 1998; for other genes please see text). Pheromone receptor gene *pre-1* (on Chr 3) and pheromone gene *ccg-4* (on Chr 1) and their products regulated by *matl-1* are also shown in white. Pheromone receptor gene *pre-2* (on Chr 7) and pheromone gene *mfa-1* (on Chr 5) and their products regulated by *matl-2* are also shown in grey. The cell walls as the margins of the cells are presented by the outer thick black lines, under which the protoplast membrane is located. Expressed pheromone receptors integrate into the membrane to which the cognate pheromones must bind for pheromone signaling transfer. In each individual trichogyne-microconidium combination, the essential protein products are selectively expressed so that courtship and fertilization can take place between a female trichogyne and a male cell of the opposite mating type. Gene *fmf-1* on Chr 1 shown as grey-white striped is a *Schizosaccharomyces pombe* *Sten*-like gene for an HMG-box TF that is required in both female- and male-acting cells for regulation of pheromone signalling and fertilization. 76

- 2.4 Life cycle of a tetrapolar heterothallic Agaricomycete with nuclear phases (haploid, dikaryotic, diploid) under complex network control of development by the two *mat* loci and additional environmental factors as known in *C. cinerea*. Heterokaryosis with different alleles in both *mat* loci, *HD* and *PR*, is necessary to form a fertile dikaryon after mycelial fusion of two compatible self-sterile homokaryons. General environmental factors with influences on mycelial growth of homokaryons and dikaryon and fruiting body induction and maturation are nutrients, temperature, humidity and conditions of aeration; increase in CO₂ blocks mycelial growth and (normal) development. On the dikaryon, dark and blue light phases, in close interaction with *mat* genes of both loci, determine the onset and progress of fruiting or, dependent on colony age and the light intensity, alternatively the disintegration of a dikaryon into its homokaryotic components (e.g. in *C. cinerea* via light-induction of oidiophores in aerial mycelium and mitotic haploid oidia produced at their tips, Polak *et al.* 1997; deduced from work of *S. commune*, increased mycelial aeration interrupts *HD*-regulated nuclear pairing and subsequently *PR*-induced gene expression, Schuur *et al.* 1998). Fruiting body development at various subsequent stages is further controlled by dark/light conditions and CO₂ concentrations, karyogamy to proceed in the basidia needs blue light and compatible *PR* genes, while meiosis is blocked by nitrogen. 93
- 2.5 Mating-type loci *matHD* (in short *HD*, classically the *A* loci) and *matPR* (in short *PR*, classically the *B* loci) of *C. cinerea*. **A**. Simplified schematics of *mat* loci and their allelic interactions in formation of a fertile dikaryon from two compatible sterile homokaryons having different *mat* loci alleles. Gene in different HD and different

PR alleles are marked in white and grey boxes, respectively. A black bar within *HD1* genes and a white bar within *HD2* genes indicate the sequence regions encoding the distinct homeodomains in the TF products of the *HD* locus. *Ph* and *Ste3* designate pheromone precursor and pheromone receptor genes in the *PR* locus, respectively. The second number in names of genes symbolizes the respective alleles. Within the dikaryon, dashed lines with two-sided arrows mark for the fertile dikaryon compatible interactions between genes of the *HD* and genes of the *PR* locus. **B. Top:** The structures of the two best characterized and fully sequenced *A* alleles (*A42* and *A43*) with three to five complete or partial paralogous *HD1-HD2* gene pairs. Grey shading indicates regions of high homology between *A42* and *A43*, non-shaded regions indicate levels of high sequence dissimilarity. The bipartite *A* locus is split into the *A α* sublocus with the functional gene pair **a** and the *A β* sublocus with functional gene pairs **b** and **d** (positions and extensions of the *A α* and *A β* subloci are marked by vertical black lines above the two maps). The internal 7 kb-long region of homologous sequence in between allows free recombination between *A α* and *A β* . Positions and names of gene pairs (**a** to **e** pair) are indicated below the subloci by horizontal black and grey lines, where black means group with proven function and grey no shown function. Different alleles of gene pairs in *A42* and *A43* are characterized by different patterns in the respective gene boxes, with the arrows indicating the direction of transcription; identical patterns mark identical alleles. Combinations of *HD1* and *HD2* genes of allelic gene pairs shown by cloning and transformation to functionally interact are marked by dashed lines with two-sided arrows. Genes of conserved sequences (*mip* and *β -fg*) flanking upstream and downstream the alleles of the *HD* locus are presented by black boxes. **B. Bottom:** The structures of the two well characterized *B* alleles *B24* and *B43*. Positions and names of paralogous groups of genes (Group 1 to Group 3) are indicated by horizontal black lines, different alleles of genes are marked by different patterns in the respective gene boxes. *Ph* and *Ste3* genes of allelic groups shown by cloning and transformation to functionally interact are marked by dashed lines with two-sided arrows. Extra genes with grey bordered boxes were found by genome sequencing and are not yet characterized. Only the *B43* allele is fully sequenced making it is possible that there are additional unidentified genes in *B42*. 95

- 4.1 Representatives of white-rot (A), brown-rot (B), grass-rot (C), plant symbiotic (D), insect fungi symbiotic (E), plant parasitic (F), and insect parasitic (F) fungi. 190
- 4.2 The life cycle of a basidiomycete fungus and its nuclear pairing and fusion. A, the life cycle of a basidiomycete fungus. B, in dikaryons, karyogamy occurs at the end of the life cycle immediately before the onset of meiosis, with the two parental nuclei remaining unfused and sharing the same cytoplasm throughout the life cycle in cells. 192
- 4.3 Schematic view of the plant lignocellulose structure and enzymes involved in their degradation. Polymeric cellulose microfibrils are tightly packed in ordered layers surrounded by branched hemicellulose chains and aromatic lignin heteropolymers. 203

- 4.4 Overview of laccase expression regulation in fungal cells. This figure summarizes the factors that induce differential laccase gene expression and the mechanisms of laccase gene regulation pathways reported to date. 207
- 5.1 A di-allele matrix. The constituent homokaryons of selected heterokaryons are recovered by protoplasting. The homokaryons are crossed in all possible combinations and evaluated for the trait of interest. In this way the breeding value of each homokaryon can be estimated in different genetic backgrounds and used to make a selection the parental homokaryons to generate segregating populations. 227
- 5.2 A breeding scheme to introduce QTL of donor lines into two acceptor lines. This approach intends to avoid inbreeding depression and is described in details in the "Outline of an effective QTL analysis and breeding strategy" paragraph. All blue lines indicate outcrosses and red lines backcrosses. 230
- 6.1 Examples of Agaricomycetes species with diverse lifestyles and fruiting body morphologies. Top row from left to right three white rot species, *Flammulina velutipes*, *Nemecomyces mongolicus* and *Cyclocybe aegerita*. The middle row shows litter decomposers, *Coprinopsis pseudonivea* and *Mycena renati*, the latter being a representative of a genus with the most expanded gene coding repertoires known to date. Also shown are two ectomycorrhiza species, the puffball-shaped *Pisolithus tinctorius* and the death cap *Amanita phalloides*, the latter being a focus of intense research on toxin systems. Bottom row: *Calocera cornea*, an early-diverging mushroom-forming species in the Dacrymycetes that causes brown rot; *Hygrophoropsis aurantiaca* (Boletales), another brown rot species, and *Amanita vittadinii* an edible *Amanita* that is considered a saprotrophic litter decomposer. 241
- 6.2 The accumulation of genomes of mushroom-forming fungi over the last 2 decades (A) and the frequency of keywords in abstracts of corresponding papers illustrated as a word cloud (B). Word size corresponds to frequency of occurrence. 243
- 6.3 The distribution of assembly sizes (A) and the number of protein coding genes (B) in published genomes available in MycoCosm (as of January 2024). Data based on statistics provided by the Joint Genome Institute. Data for *Mycena olivaceomarginata* (97000 genes) have been omitted from (B) for visualization purposes. 255
- 8.1 Mechanism of the CRISPR/Cas9 gene editing system. 308
- 9.1 Mutation accumulation studies are designed to estimation mutation rates by growing cells in the absence of selection. A) The fluctuation assay was used originally to show that mutations are random and not caused by the selective environment (Luria-Delbrück experiment). In this example, mutation at the *ura3* locus is used to estimate rates of mutation to null alleles. A single genotype is grown in permissive conditions that counter selects for mutations, in this case minimal medium lacking uracil. After growing numerous small populations in numerous replicates, some populations are used to estimate total population per well (e.g., via a Coulter counter), and other populations are plated in their entirety onto selective media (e.g., 5FOA) to estimate the total number of *ura-* cells per population. B) Mutation accumulation with yeast involves streaking single colonies out from plate to plate. Using the estimated num-

ber of cells per colony allows the number of transfers to be translated to the number of generations. Transfers can be streaks or using spread plating of diluted colonies. C) With a mold species such as *Aspergillus*, an individual clone from a spore can be isolated and used to establish lines that are independent. After a certain number of days, the margin of the mycelium can be sampled for conidia, and a small number of conidia used to start the next passage after dilution in water. D) For species that do not sporulate, mycelium can be transferred in between passages. In order to reduce the effective population size Bezemova et al. (2020) used thin tubes of either 4 mm or 0.8 mm thickness to reduce competition among nuclei at the leading margin of the growing mycelium. After the mycelium reaches the end of the tube, the hyphae are seeded into clean tubes of media for another passage. 330

- 9.2 Genetic diversity in cultivated *A. bisporus* strains is limited. Phylogeny of major clades of *Agaricus bisporus* and related species based on mitochondrial DNA SNPs. Cultivars and their escapes are only known from Clades Europe II & III, which show low diversity within each clade. Shown in pie charts are the proportion of strains assigned to cultivar, wild, or escape. Branch lengths are scaled to millions of years using a molecular clock. 333
- 9.3 Evidence of strain degeneration in *Cordyceps militaris*. A-B. Healthy fruiting body production observed on typical strain grown on wheat (A) and vigorous fruiting on an inoculated pupa (B). C-D. Degenerated strains after repeated subculturing and produce fewer and poorly pigmented fruiting bodies in jar cultures (C) and fail to produce fruiting bodies when inoculated on a pupa (D). Photos generously provided by Mengqian Liu and Caihong Dong, Chinese Academy of Sciences. 335
- 10.1 Mass of tissue symptom of Dry Bubble Disease 353
- 10.2 Split stem symptom of Dry Bubble Disease 354
- 10.3 *Trichoderma aggressivum* f. *aggressivum* (Taz) growing on the surface of the casing. It is dark green with a white fringe. 357
- 10.4 a) White fluffy growth symptom of cobweb disease and b) Fleshy color spotting symptom 360
- 10.5 *S. megalocarpus* sporangiospores on a branched sporangium 362
- 10.6 Zygosporangia of *S. megalocarpus* that show the dark colored, thick-walled spores 363
- 10.7 *S. megalocarpus* growing on an *A. bisporus* with the web or bread hair-like growth 363
- 10.8 Internal decay of the mushroom fruiting body caused by *S. megalocarpus* 364
- 10.9 Wet Bubble Disease showing the amber colored drops symptom 365
- 10.10 Bacterial Blotch discoloration symptom 367
- 10.11 Misshapen mushrooms symptom of mummy disease 370
- 10.12 a) Premature cap opening symptom of LaFrance Disease b) Drumstick-shaped mushroom symptom 373
- 10.13 Symptoms of MVX-PD: bare patches in crop 375
- 10.14 Symptoms of MVX: coffee-colored mushrooms 376
- 10.15 Mushroom phorid fly (*Megaselia halterata*) life cycle 378

- 10.16 *S. carpocapsae* nematodes at different life stages. A) *S. carpocapsae* infective juveniles. B) Mature adult *S. carpocapsae* bursting from a dead *M. halterata* larva. C) *S. carpocapsae* within a larval cadaver. 380
- 10.17 Stratiolaelaps scimitus predatory mites. A) A *S. scimitus* mite (PM) next to a *Tyrophagus* spp. mold mite with a penny for scale. B) Adult *S. scimitus* mites feeding on a *M. halterata* larva. 381
- 10.18 Attract-and-kill station on a farm. This attract-and-kill station used natural light as an attractant at a vent window, with an electrostatically charged net blocking fly movement. Powdered insecticides were applied to the insecticidal nets to kill adults that were behaviorally trapped at the station. 382
- 10.19 Fungus gnat (*Lycoriella ingenua*) life cycle 383
- 11.1 Examples of white and cremini button mushrooms in growth rooms (A and B), and all three types from markets (C: top, portobello; middle, cremini; bottom, white). 397
- 11.2 Full colonization of the substrate by *A. bisporus* mycelia 402
- 11.3 Mushroom picking is a labour-intensive process that requires significant hand skills to be effective 406
- 11.4 Packaged whole and sliced button mushrooms as well as portobello mushrooms 407
- 12.1 Substrate preparation for growing oyster mushrooms 423
- 12.2 Bagging and sterilization of substrates for growing oyster mushrooms 424
- 12.3 Incubation of fruiting substrates inoculated with oyster strains 425
- 12.4 Fruiting of oyster mushrooms 426
- 13.1 Fruiting body of *L. edodes* 433
- 13.2 Mixing sawdust and other materials for growing *Lentinula edodes* 439
- 13.3 Bagging of mixed substrate for growing *Lentinula edodes* 440
- 13.4 Traditional method for sterilizing shiitake-growing substrates 441
- 13.5 Inoculation of Shiitake spawns into sterilized substrate 442
- 13.6 Mycelium growth and colonization of substrates by Shiitake strains 443
- 13.7 Formation of dark brown mycelium 443
- 13.8 Harvesting shiitake grown on inclined artificial logs 445
- 13.9 Injecting water into the bag 446
- 14.1 Life cycle of the straw mushroom 453
- 14.2 Cultivation equipment and facility for the straw mushroom. A) straw mushroom cultivation room; B) the corridor equipment of straw mushroom cultivation room; C) Straw mushroom cultivation shelf; D) Cultivated straw mushrooms 458
- 15.1 Life cycle of *Flammulina filiformis* 467
- 15.2 Flow chart of *Flammulina filiformis* cultivation technology 472
- 15.3 Wrapping with plastic sheet 478
- 16.1 Seedlings of *Pinus armandii* before addition of truffle spore inoculum 488
- 16.2 Differences in the growth of seedlings of *Pinus armandii* inoculated with truffle spores (right) or not (left). 489
- 16.3 Ectomycorrhizas formed by *Pinus armandii* and *T. indicum*. a: naked eye view of mycorrhizas under one root system; b: one mycorrhiza under dissecting microscope 489

- 16.4 Measuring the percentage of truffle mycorrhizal colonization 489
- 17.1 *Auricularia heimuer* sexual life cycle 503
- 17.2 Representative cultivars of *Auricularia cornea* 507
- 17.3 Small-scale (left) and large-scale (right) autoclave 509
- 17.4 Manual (left) and automated (right) spawn inoculations. 510
- 17.5 A: Incubation room components and set up. B: Full mycelia colonization. 511
- 17.6 Bag openings for inducing fruiting body formation after full colonization 512
- 17.7 Hole-puncher for fungal fruiting bags 513
- 18.1 Production facility of *P. portentosus* in Jinghong City of Yunnan Province, China 521
- 18.2 *P. portentosus* produced in large-scale factory facility. 526
- 18.3 Appearance of Cultivar YL1701-2 530
- 19.1 Fruiting body of *Hericium erinaceus* 541
- 19.2 Fruiting room for *Hericium erinaceus* 548
- 20.1 Tree cutting and wood sectioning for growing Ganoderma mushrooms 561
- 20.2 Bagging of wood sections for growing Ganoderma mushrooms 562
- 20.3 Stacking and sterilizing bagged wood segments for growing Ganoderma mushrooms 563
- 20.4 Arrangement, bag removal, and soil covering of wood sections colonized with Lingzhi mycelia 564
- 20.5 Growth management of Ganoderma fruiting bodies 567
- 20.6 Collection of Ganoderma spore powder using paper bag sleeves 568
- 20.7 Collection of Ganoderma spore powder in a nonwoven shed 536
- 20.8 Management of Lingzhi production in shelf-based cultivation of substitute substrates 571
- 20.9 Root art and auspicious cloud Lingzhi 572
- 20.10 Cultivation of large Lingzhi fruiting bodies 573
- 20.11 Cultivation of antler-shaped ganoderma 573
- 20.12 Cultivation of tower-shaped bonsai Lingzhi 574
- 21.1 *Cyclocybe chaxingu* (HFJAU1332). A, B: Basidiomata; C: Basidia; D: Pleurocystidia; E: Basidiospores; F: Pileocystidia; G: Cheilocystidia; H: Colonies. Bars=10 μm. 579
- 21.2 Differences in pileus color exhibited by different cultivars. a, *Gancha AS-1*; b, *Gucha 1*; c, *Gucha 2*; d, A cultivar named *Baicha* with white pileus 584
- 21.3 The mycelium of *Cyclocybe chaxingu* growing inside mushroom bags 589
- 21.4 Wall bag cultivation of *Cyclocybe chaxingu* 589
- 22.1 Four cultivated cordycipitoid fungal species from natural habitat. A, *Ophiocordyceps sinensis*; B, *Cordyceps militaris*; C, *Tolypocladium guangdongense*; D, *Cordyceps chanhua*. 597
- 22.2 Fruiting body cultivation of *Ophiocordyceps sinensis* on host insects 601
- 22.3 Fruiting body cultivation of *Cordyceps militaris* on artificial media (left) and insects (right) 602
- 22.4 Fruiting body cultivation of *Tolypocladium guangdongense* on artificial media 603
- 22.5 Fruiting body cultivation of *Cordyceps chanhua* on artificial media 604

- 23.1 A: Maximum likelihood (ML) phylogenetic analysis of species in the genus *Morchella* based on a four-gene data set (ITS plus *EF1-a* plus *RPB1* plus *RPB2*), with *Verpa* and *Disciotis* as outgroups. B: The relationships among the proposed four groups (blushing morels, yellow morels, black morels, and semi-free-capped morels), three clades (*Rufobrunnea* Clade, *Esculenta* Clade, and *Elata* Clade), and three sections (*Rufobrunnea*, *Morchella*, and *Distantes*). 611
- 23.2 The revised life cycle of *Morchella* based on Du and Yang (2021) 616
- 23.3 Main cultivars. A: *Morchella importuna*; B: *Morchella sextelata*; C: *Morchella eximia* 618
- 23.4 Cultivation stepwise protocols and cultivation modes of true morels 620
- 23.5 Harvesting and processing of cultivated true morels 625
- 24.1 Examples of mushroom stamps 641
- 24.2 Local merchandise based on large underground mushroom mycelia reported in scientific and popular press 642
- 24.3 Underground mycelial network of *Termitomyces* 643
- 24.4 Harvested fruiting bodies of *Tricholoma matsutake* 643
- 24.5 Commercially cultivated and harvested *Flammulina filiformis* and naturally found fruiting bodies 644
- 24.6 Representative pictures of *Ophiocordyceps sinensis* 646
- 24.7 *Ganoderma* in building wall and porcelain 652
- 24.8 Tianquan Mushroom Museum in Changshan, Zhejiang, China 653
- 24.9 Representative mushroom dishes 654
- 25.1 Biosynthesis pathway of terpenoid in Basidiomycetes and Ascomycetes. This schematic pathway was drawn based on the information reviewed in reference 662
- 25.2 Shikimate biosynthetic pathway in mushroom. The full name and the respective abbreviation of compounds are as follows, 5-enolpyruvylshikimate-3-phosphate (EPSP); phosphoenolpyruvate (PEP); 3-deoxy-D-arabino-heptulosonate-7-phosphate (DAHP); dehydroquinate (DHQ); 3-dehydroshikimate (DHS); shikimate-3-phosphate (S-3-P); The bullets/numbering (1)–(7) indicate respective enzymes catalyzing the seven steps of shikimate pathway: (1) DAHP synthase; (2) dehydroquinate synthase (DHQS); (3) dehydroquinate dehydratase (DHQase); (4) Shikimate dehydrogenase (SD); (5) Shikimate kinase (SK); (6) EPSP synthase; (7) chorismate synthase. 665
- 27.1 Structures of toxic cyclopeptides 740
- 28.1 Structures of C8-oxylipins, their perceived odour and biosynthetic pathway of oct-1-en-3-ol synthesis via the oxygenation of linoleic acid to 10-(S)-hydroperoxy-9,11-octadecadienoic acid (10-HPOD) by a dioxygenase (DOX) and the subsequent cleavage by a hydroperoxide lyase (HPL). 757
- 28.2 Schematic biosynthesis of isopentenyl diphosphate (IPP) and dimethylallyl diphosphate (DMAPP) in fungi, which are linked to generate the geranyl diphosphate (GPP), farnesyl diphosphate (FPP) and geranylgeranyl diphosphate (GGPP), the precursors for volatile and semi-volatile terpenes. The enzymes in the boxes are acetyl-CoA C-acetyltransferase (ACAT), 3-hydroxy-3-methylglutaryl-CoA synthase

- (HMGS), 3-hydroxy-3-methylglutaryl-CoA reductase (HMGR), mevalonate kinase (MVK), phosphomevalonate kinase (PMVK), mevalonate diphosphate decarboxylase (MVD), isopentenyl diphosphate isomerase (IDI), GPP synthase (GPPS), FPP synthase (FPPS), GGPP synthase (GGPPS). 762
- 28.3 Structures of several volatile terpenes found in mushrooms. Beneath the trivial names are listed the perceived odours, if described. 766
- 28.4 Structures of volatile sulphur compounds (VSC) found in mushrooms 770
- 29.1 Lion's mane (*Hericium erinaceus*) growing from a kit indoors 792
- 29.2 Blue Oyster Mushrooms (*Pleurotus ostreatus*) ready for harvest, growing under 6500K LED lights. 792
- 29.3 Fruiting mushroom blocks on a shelf at Nature Lion's mushroom farm 793
- 29.4 Harvesting lion's mane mushrooms. (*Hericium erinaceus*) 793
- 29.5 Adding chopped straw to a sack and pasteurize in boiling water. 798
- 30.1 The real mycelium-based leather products of Reishi™ from MycoWorks and Mylea™ from Mycotech Lab. 821
- 30.2 The real mycelium-based packing products from Grown.bio 822
- 30.3 The real mycelium-based building products from Mogu 823
- 30.4 The real mycelium-based furniture interior design products from Grown.bio and Ecovative 823
- 30.5 The mycelium-based elect electronic components. (a) mycelium skin with Cu-Au traces; (b) demonstration photograph of a mycelium-based sensor board with different elements; (c) demonstration of the mycelium-based battery; (d) section view of the mycelium zinc-carbon battery. 824
- 30.6 Commercial-scale production of mycelium-based materials. (a) Bioreactor of mycelium leather production at VTT, (b) vertical farming of aerial mycelium foam ©Bolt Threads, (c) and (d) commercial-scale production farm of ©Ecovative 828

Tables

- 1.1 Jaccard similarity among six continents 9
- A1 The classification of edible and medicinal mushrooms above family level, primarily sourced from the Mycobank database along with additional references. 12
- 2.1 Molecular structure of mat loci of cultivated mushroom species from the Basidiomycota, Agaricomycetes 101
- 3.1 Common cultivated mushrooms and their germplasm preservation centers in the world 158
- 3.2 Representative examples of population genetic studies of edible mushrooms 177
- 5.1 An overview from all genetic linkage maps generated for edible mushrooms since 1993 218
- 5.2 Quantitative trait loci (QTL) studies in edible mushrooms done from 1999 up to 2023 220

- 7.1 Advances in research on edible fungi transformation in the past two decades 284
- 7.2 Examples of successful gene editing and silencing in edible fungi 287
- 10.1 Prevalence of two viral diseases on *Agaricus bisporus* mushrooms 372
- 11.1 Compositions of two common types of composts used for growing *A. bisporus* 401
- 16.1 Advantages and disadvantages of three different types of inoculums in producing mycorrhizal seedlings 487
- 20.1 Commonly used formulas for Lingzhi substitute substrate cultivation 570
- 22.1 Taxonomy and host range of four cultivated cordycipitoid fungi 597
- 25.1 Representative mushroom pigments sorted by their biosynthesis pathways and the mushroom producers 669
- 26.1 Effects of adding edible mushrooms on the rheological, cooking, and microstructural properties of staple composite foods 714
- 26.2 Selected dietary supplements based on mushrooms 717
- 26.3 Cosmetic products containing mushrooms and their ingredients 722
- 27.1 The new species of poisonous mushrooms reported from 2012 to 2022 737
- 28.1 Key aroma compounds of some of the most valuable mushrooms 774
- 29.1 A list of select mushroom grain spawn and grow kit suppliers 795
- 29.2 Hardwood sawdust & wheat bran growing block recipe 797
- 29.2 Straw growing block recipe 797
- 30.1 The spent mushroom substrates (SMS) used for organic fertilizer productions 811
- 30.2 The spent mushroom substrates (SMS) used for new mushroom cultivations 813
- 30.3. The spent mushroom substrates (SMS) used for biofuel productions 815
- 30.4 Examples of spent mushroom substrates (SMS) used for crop disease and pest management 816
- 30.5 Bioremediation of environmental contaminants by spent mushroom substrates (SMS) 818
- 30.6 Mushroom species used for mycelium-based materials 826